

**PDC2017
Tokyo, Japan**

IAA-PDC-17-03-P07

Please send your abstract to iaapdc (at) iaamail.org

You may visit www.pdc.iaaweb.org

*(please choose one box to be checked)
(you may also add a general comment - see end of the page)*

- Key International and Political Developments**
- Advancements and Progress in NEO Discovery**
- NEO Characterization Results**
- Deflection and Disruption Models & Testing**
- Mission & Campaign Designs**
- Impact Consequences**
- Disaster Response**
- Decision to Act**
- Public Education & Communication**

**OBSERVATION OF NEAR-EARTH OBJECT (1566) ICARUS AND THE SPLIT
CANDIDATE 2007 MK₆**

**S. Urakawa^{(1)*}, K. Ohtsuka⁽²⁾, S. Abe⁽³⁾, D. Kinoshita⁽⁴⁾, H. Hanayama⁽⁵⁾,
T. Miyaji⁽⁵⁾, S. Okumura⁽¹⁾, K. Ayani⁽⁶⁾, S. Maeno⁽⁶⁾, D. Kuroda⁽⁷⁾, A. Fukui⁽⁷⁾,
N. Narita⁽⁸⁾⁽⁹⁾⁽¹⁰⁾, G. L. Hashimoto⁽¹¹⁾, Y. Sakurai⁽¹¹⁾, S. Nakamura⁽¹¹⁾, J.
Takahashi⁽¹²⁾, T. Tanigawa⁽¹³⁾, O. Burhonov⁽¹⁴⁾, K. Ergashev⁽¹⁴⁾, T. Ito⁽¹⁰⁾,
F. Yoshida⁽¹⁰⁾, M. Watanabe⁽¹⁵⁾, M. Imai⁽¹⁶⁾, K. Kuramoto⁽¹⁶⁾, T. Sekiguchi⁽¹⁷⁾,
and M. Ishiguro⁽¹⁸⁾**

⁽¹⁾Japan Spaceguard Association, 1716-3 Okura, Bisei, Ibara, Okayama 714-1411, Japan, ⁽²⁾Tokyo Meteor Network, 1-27-5 Daisawa, Setagaya-ku, Tokyo 155-0032, Japan, ⁽³⁾Nihon University, 7-24-1 Narashinodai, Funabashi, Chiba 274-8501, Japan, ⁽⁴⁾National Central University, 300 Jhongda Rd., Jhongli, Taoyuan 32001, Taiwan, ⁽⁵⁾Ishigakijima Astronomical Observatory, National Astronomical Observatory of Japan, 1024-1 Arakawa, Ishigaki, Okinawa 907-0024, Japan, ⁽⁶⁾Bisei Astronomical Observatory, 1723-70 Okura, Bisei, Ibara, Okayama 714-1411, Japan, ⁽⁷⁾Okayama Astrophysical Observatory, National Astronomical Observatory of Japan, Asakuchi, Okayama 719-0232, Japan, ⁽⁸⁾Astrobiology Center, 2-21-1 Osawa, Mitaka, Tokyo, 181-8588, Japan, ⁽⁹⁾The University of Tokyo, 7-3-1 Bunkyo-ku, Tokyo 113-0033, Japan, ⁽¹⁰⁾National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo, 181-8588, Japan, ⁽¹¹⁾Okayama University, 3-1-1 Tsushimanaka, Kita-ku, Okayama 700-8530, Japan, ⁽¹²⁾University of Hyogo, Sayo, Hyogo 679-5313, Japan, ⁽¹³⁾Sanda Shoukan Highschool, 101 Gakuen, Sanda, Hyogo, 669-1337, Japan, ⁽¹⁴⁾Ulugh Beg Astronomical Institute Uzbekistan Academy of Sciences, 33 Astronomicheskaya str., Tashkent, 100052, Uzbekistan, ⁽¹⁵⁾Okayama University of Science, 1-1 Ridai, Kita-ku, Okayama 700-0005, Japan, ⁽¹⁶⁾Hokkaido University, Kita 8, Nishi5, Kita-ku, Sapporo, Hokkaido 060-0810, Japan, ⁽¹⁷⁾Hokkaido University of

Education, Hokumon 9, Asahikawa, Hokkaido 070-8621, Japan, ⁽¹⁸⁾Seoul National University, Gwanak, Seoul 151-742, Korea

Keywords: Asteroids, Taxonomy, Rotation, Photometry

ABSTRACT

Background & Aim: A numerical simulation proposes that the origin of near-Earth object 2007 MK₆ (hereafter, MK6) is a near-Earth object (1566) Icarus (hereafter, Icarus) [1]. In addition to it, the orbital parameters of the daytime Taurid-Perseid meteor swarm are in good agreement with those of Icarus. Thus, it is considered that MK6 is split from the parent object Icarus by a rotational fission and/or an impact event, and the produced dust became to the daytime Taurid-Perseid meteor swarm. To confirm such a hypothesis, we need to obtain the observational evidence that the color indices of Icarus and MK6 are same. Moreover, if MK6 split by the rotational fission due to the YORP effect, the rotation period of Icarus would shorten compared with the past rotation period. When the MK6 split by an impact event, the rotation period of MK6 would shorter than the spin limit of 2.2 hours. We require the observation for Icarus and MK6 to test these hypotheses.

Observations: We conducted the observations for Icarus in 2015 and MK6 in 2016, respectively. The observation summary is shown as followings: Icarus (June 2015): Nayoro Observatory 1.6 m Pirka telescope of the Hokkaido University (visible photometry), Ishigakijima Astronomical Observatory (IAO) 1.05 m Murikabushi telescope (g' , R_C , and I_C band simultaneous photometry), Maidanak Observatory (MO) 0.6 m telescope (R band photometry), Nishi-Harima Astronomical Observatory (NHAO) Nayuta 2.0m telescope (J , H , and K_S band simultaneous photometry), Lulin and Kinmen Observatory 0.4 m telescopes (visible photometry), Lowell Observatory (LO) 1.1 m, 1.8 m, and 4.3 m telescopes (visible photometry and visible spectroscopy), MK6 (June - July 2016): IAO 1.05 m Murikabushi telescope (g' , R_C , and I_C band simultaneous photometry), MO 1.5 m telescope (R band photometry), NHAO Nayuta 2.0 m telescope (J , H , and K_S band simultaneous photometry), Lulin Observatory 1.0 m telescope (visible photometry), LO 1.8 m and 4.3 m telescopes (visible photometry and visible spectroscopy), University of Hawaii 2.24 m telescope (visible photometry), Okayama Astrophysical Observatory 1.88 m telescope (g' , r' , and z' band simultaneous photometry).

Results: Previous studies indicated that the taxonomy of Icarus is an S-type or a Q-type [2][3]. We obtained that the color indices $g' - R_C$, $R_C - I_C$, $J - H$, and $H - K_S$ are 0.828 ± 0.027 mag, 0.397 ± 0.025 mag, 0.515 ± 0.048 mag, and 0.069 ± 0.030 mag, respectively. These are consistent with the color indices of an S-type asteroid. Moreover, the color indices implied the slight rotational color variation, though the further data analysis is needed. On the other hand, the color indices and the rotation period of MK6 have not been revealed in the previous study. In addition to the color indices of Icarus, we will present the result the color indices of MK6 and the rotation period of both Icarus and MK6.

References: [1] Ohtsuka K. et al. (2007) *ApJ*, 668, L71-L74. [2]Chapman C. R., Morrison D., and Zellner B. (1975) *Icarus*, 25, 104-139. [3]Hicks M. D., Fink U., and Grundy W. M. (1998) *Icarus*, 133, 69-78.

* Corresponding author

Email address: urakawa@spaceguard.or.jp