

**PDC2019  
Washington, DC, USA**

*Please send your abstract to iaapdc (at) iaamail.org*

*You may visit [www.pdc.iaaweb.org](http://www.pdc.iaaweb.org)*

*(please choose one box to be checked)  
(you may also add a general comment - see end of the page)*

- Key International and Political Developments**
- Advancements and Progress in NEO Discovery**
- NEO Characterization Results**
- Deflection and Disruption Models & Testing**
- Mission & Campaign Designs**
- Impact Consequences**
- Disaster Response**
- Decision to Act**
- Public Education & Communication**

**Asteroid to Airburst; Comparing Semi-analytical Airburst Models to Hydrocodes**

**Sarah McMullan<sup>(1)</sup> and Gareth Collins<sup>(2)</sup>**

*<sup>(1)(2)</sup> Impact and Astromaterials Research Centre, Dept. Earth Science and Eng.,  
Imperial College London, SW7 2BP, UK, (+44)7960109673,  
[s.mcmullan16@imperial.ac.uk](mailto:s.mcmullan16@imperial.ac.uk)*

**Keywords:** *Numerical Simulations, Airbursts, Energy Deposition*

**ABSTRACT**

Near-Earth Objects, 1-100 m in size, are abundant, difficult to observe, and strike the Earth with little to no warning [1]. There is likely to be at least one significant event in a human lifespan (e.g. Chelyabinsk, Tunguska), so being able to model their potential energy deposition and damage on the ground accurately is imperative. Semi-analytical models are a fast predictors of energy deposition, but they make many assumptions and have several free parameters that are poorly defined [2-4]. These parameters are usually tuned to fit a particular data set, such as the 2013 Chelyabinsk event in Russia [4]. In previous work, we showed that it is possible to tune multiple different semi-analytical models to match the Chelyabinsk energy deposition [5], however, when these same models are then applied to a different event, such as Tunguska, the results diverge [6]. Moreover, all of the existing semi-analytical models in popular use predict unphysically large spreading limits, several times greater than the initial asteroid diameter, which raises concerns about their predictive capacity.

Shock physics codes rely on fewer simplifying assumptions, but have a higher computational cost than semi-analytical models. Here we use the iSALE shock physics code [7-9] to simulate the atmospheric entry of meteoroids. We consider the effect of asteroid strength and porosity on the rate of deceleration and spreading. From our simulation results, we produce synthetic energy deposition curves with known initial entry conditions. This provides an additional data set with which to calibrate existing semi-analytical models and develop new approaches for the fast prediction of airburst outcomes necessary for probabilistic hazard assessment or rapid decision making.

### References:

- [1] Harris, A. W. Boslough, M., Chapman, C. R., Drube, L., Michel, P., & Harris, A. W. (2015). Asteroid Impacts and Modern Civilization: Can We Prevent a Catastrophe? In *Asteroids IV* (pp. 835-854) Univ. of Arizona, Tuscon.
- [2] Chyba, C. F., Thomas, P. J., & Zahnle, K. J. (1993). The 1908 Tunguska explosion: atmospheric disruption of a stony asteroid. *Nature*, 361, 40-44.
- [3] Hills, J. G., & Goda, M. P. (1993). The Fragmentation of Small Asteroids in the Atmosphere. *The Astronomical Journal*, 105, 1114-1144.
- [4] Avramenko, M. I., Glazyrin, I. V., Ionov, G. V., & Karpeev, A. V. (2014). Simulations of the airwave caused by the Chelyabinsk superbolide. *Journal of Geophysical Research: Atmospheres*, 119, 7035-7050.
- [5] Brown, P.G., & 32 co-authors (2013). A 500-kiloton airburst over Chelyabinsk and an enhanced hazard from small impactors. *Nature*, 503, 238-241.
- [6] McMullan, S., & Collins, G. S. (2019). Uncertainty Quantification in Continuous Fragmentation Airburst Models. *Icarus (in review)*.
- [7] Amsden, A., Ruppel, H., and Hirt, C. (1980). SALE: A simplified ALE computer program for fluid flow at all speeds. *Los Alamos National Laboratories Report, LA-8095:101p*. Los Alamos, New Mexico: LANL.
- [8] Collins, G. S., Melosh, H. J., and Ivanov, B. A. (2004). Modeling damage and deformation in impact simulations. *Meteoritics and Planetary Science*, 39, 217--231.
- [9] Wünnemann, K., Collins, G., and Melosh, H. (2006). A strain-based porosity model for use in hydrocode simulations of impacts and implications for transient crater growth in porous targets. *Icarus*, 180, 514--527.

\*\*\*\*\*